

Hepatorenal syndrome: pathophysiology, diagnosis, and management

Douglas A Simonetto,¹ Pere Gines,² Patrick S Kamath¹



¹Division of Gastroenterology and Hepatology, Mayo Clinic College of Medicine and Science, Rochester, MN 55905, USA

²Liver Unit, Hospital Clinic, University of Barcelona IDIBAPS – CIBERehD, Barcelona, Spain

Correspondence to: P S Kamath
Kamath.patrick@mayo.edu

Cite this as: *BMJ* 2020;370:m2687
<http://dx.doi.org/10.1136/bmj.m2687>

Series explanation: State of the Art Reviews are commissioned on the basis of their relevance to academics and specialists in the US and internationally. For this reason they are written predominantly by US authors.

ABSTRACT

Hepatorenal syndrome (HRS), the extreme manifestation of renal impairment in patients with cirrhosis, is characterized by reduction in renal blood flow and glomerular filtration rate. Hepatorenal syndrome is diagnosed when kidney function is reduced but evidence of intrinsic kidney disease, such as hematuria, proteinuria, or abnormal kidney ultrasonography, is absent. Unlike other causes of acute kidney injury (AKI), hepatorenal syndrome results from functional changes in the renal circulation and is potentially reversible with liver transplantation or vasoconstrictor drugs. Two forms of hepatorenal syndrome are recognized depending on the acuity and progression of kidney injury. The first represents an acute impairment of kidney function, HRS-AKI, whereas the second represents a more chronic kidney dysfunction, HRS-CKD (chronic kidney disease). In this review, we provide critical insight into the definition, pathophysiology, diagnosis, and management of hepatorenal syndrome.

Introduction

Hepatorenal syndrome is a serious complication of cirrhosis that is associated with high morbidity and mortality. It is characterized by functional circulatory changes in the kidneys that overpower physiologic compensatory mechanisms and lead to reduced glomerular filtration rate. Re-establishment of adequate renal blood flow leads to improvement in renal function and is achieved with liver transplantation or vasoconstrictor drugs. The terminology, definition, and classification of hepatorenal syndrome have evolved considerably in the past 10 years owing to changes proposed for the diagnosis and staging of acute kidney injury (AKI). In this state of the art review, we focus on recent advances that have shaped the contemporary definition, diagnosis, and management of hepatorenal syndrome.

Sources and selection criteria

We searched PubMed from January 2000 to October 2019 using the keyword “hepatorenal syndrome”. We prioritized randomized clinical trials (RCTs), systematic reviews, and meta-analysis, when available, excluding case reports and case series. We also reviewed published management guidelines from websites of professional societies (American Association for the Study of Liver Diseases; European Association for the Study of the Liver; Kidney Disease: Improving Global Outcomes; and others). We included only full length, peer reviewed studies published in English, with the exception of recent high profile RCTs available only in abstract form.

Epidemiology and definitions

Acute impairment of renal function, as determined by an increase in serum creatinine, is reported in 19-26% of patients admitted to hospital with cirrhosis and 32% of patients with severe alcohol associated hepatitis.¹⁻⁴ Serum creatinine, however, underestimates decline in renal function in patients with cirrhosis owing to impaired hepatic production of creatine (the precursor of creatinine), reduced muscle mass, tubular secretion of creatinine, and inaccurate measurement of creatinine by calorimetric methods in the presence of elevated serum bilirubin.^{5,6} Therefore, alternative biomarkers with greater accuracy in patients with cirrhosis are urgently needed. Incorporating cystatin C, a cysteine proteinase inhibitor, into conventional creatinine based glomerular filtration rate (GFR) equations results in better diagnostic performance compared with conventional equations alone.⁷

Despite the limitations, serum creatinine remains the most widely available and inexpensive assay for estimation of GFR. In a non-steady state such as acute renal failure, serial changes in creatinine and urine output better reflect renal function. Consequently, the definition of acute renal failure has evolved over the past two decades. The RIFLE (risk, injury, failure, loss, and end stage kidney disease) classification, generated at the Consensus Conference of the Acute Dialysis Quality Initiative Group in 2002, consisted of percentage changes in serum creatinine or GFR, a decrease in urine output, or both.⁸ This classification was later refined by a second multidisciplinary collaborative forum held in 2005, the Acute Kidney



SHOUT
SHARE IT

Injury Network (AKIN).⁹ The AKIN first proposed the term acute kidney injury (AKI) to include the entire spectrum of acute renal failure and also added an absolute increase in serum creatinine of 0.3 mg/dL within 48 hours of baseline as part of the definition of AKI. This change was based on accumulating evidence that even small acute increments in serum creatinine were associated with significant short term morbidity and mortality.^{10 11} In 2012 the Kidney Disease Improving Global Outcome (KDIGO) organization published clinical practice guidelines further refining the diagnosis of AKI on the basis of the RIFLE and AKIN criteria. The KDIGO defined AKI as an increase in serum creatinine by at least 0.3 mg/dL (26.5 µmol/L) within 48 hours, an increase in serum creatinine to at least 1.5 times baseline within the previous seven days, or urine volume below 0.5 mL/kg/h for six hours.¹²

The definition of hepatorenal syndrome has likewise evolved in the past two decades to align with the changes proposed by the RIFLE, AKIN, and KDIGO guidelines. In 1990 the International Club of Ascites (ICA) defined acute renal failure in cirrhosis as an increase in serum creatinine of at least 50% from baseline to a final concentration of at least 1.5 mg/dL. Hepatorenal syndrome was further subclassified into two clinical types: type 1, defined as rapid reduction of renal function by doubling of initial serum creatinine to a concentration of at least 2.5 mg/dL or a 50% reduction in less than two weeks in the initial 24 hour creatinine clearance to below 20 mL/min; and type 2, in which renal failure progression did not meet the criteria for type 1.¹³

Several recent studies showed that the diagnosis of AKI in patients with cirrhosis, based on an absolute increase in serum creatinine by at least 0.3 mg/dL or 50% from baseline, leads to earlier identification of patients at increased risk of longer hospital stay, multi-organ failure, admission to intensive care, and in-hospital mortality as well as 90 day mortality.¹⁴⁻¹⁸ This led the ICA to issue a revised set of consensus recommendations in 2015, incorporating a new definition and classification of AKI with modifications (box 1).¹⁹ A serum creatinine concentration obtained in the previous three months can be used as baseline when a concentration in the previous seven days is not available. The ICA also eliminated urine output in the revised definition of AKI, owing to expected lower urine output at baseline in patients with cirrhosis due to avid sodium and water retention. However, a recent study showed that patients in the intensive care unit with reduction in urine output to below 0.5 mL/kg for at least six hours had a higher mortality than did those who met only the creatinine criteria for AKI.²⁰ Therefore, accurate urine output (for example, when obtained through a urinary catheter) may be used for diagnosis and staging of AKI.²¹

The ICA also updated the definition of hepatorenal syndrome (HRS) type 1, now termed HRS-AKI. The two week interval required for doubling in serum creatinine in the previous definition could lead

Box 1: Stages of acute kidney injury according to the International Club of Ascites¹⁹

Stage 1

Increase in serum creatinine ≥ 0.3 mg/dL (26.5 µmol/L) or increase in serum creatinine ≥ 1.5 -fold to twofold from baseline

- Stage 1a
- Creatinine < 1.5 mg/dL
- Stage 1b
- Creatinine ≥ 1.5 mg/dL

Stage 2

Increase in serum creatinine at least twofold to threefold from baseline

Stage 3

Increase in serum creatinine at least threefold from baseline or serum creatinine ≥ 4.0 mg/dL (353.6 µmol/L) with an acute increase ≥ 0.3 mg/dL (26.5 µmol/L) or initiation of renal replacement therapy

to a delay in starting treatment for hepatorenal syndrome; several studies have shown that the higher the creatinine at the start of treatment, the lower the probability of reversal of hepatorenal syndrome.^{22 23} Therefore, no minimum creatinine value is needed for the diagnosis of HRS-AKI according to the updated definition. That is, HRS-AKI can be diagnosed even when the serum creatinine is below 2.5 mg/dL.

Functional kidney injury in patients with cirrhosis that does not meet the criteria for HRS-AKI is termed HRS-NAKI (that is, non-AKI) and is defined by estimated glomerular filtration rate (eGFR) rather than serum creatinine.²⁴ NAKI is divided into HRS-acute kidney disease (HRS-AKD) if the eGFR is less than 60 mL/min/1.73 m² for less than three months and HRS-chronic kidney disease (HRS-CKD) if it is less than this for more than three months (fig 1).

Pathophysiology

Our understanding of the pathophysiology of hepatorenal syndrome is mostly based on observational studies in humans because of the lack of a reproducible experimental model of hepatorenal syndrome. The challenge faced with animal models is the inability to induce severe liver injury without direct kidney toxicity as with carbon tetrachloride and thioacetamide. Nevertheless, clinical and histopathologic observations point to an uncompensated hyperdynamic circulation as the hallmark of AKI-HRS. Additional contributors to development of AKI-HRS include systemic inflammation, cirrhotic cardiomyopathy, and adrenal insufficiency.

Circulatory dysfunction

The hemodynamic changes in patients with cirrhosis are linked to sodium retention, development of ascites, and subsequent renal dysfunction.²⁵ Cirrhosis results in elevated intrahepatic vascular resistance but splanchnic vasodilatation due to

OLD NAME	NEW NAME
HRS type 1 ■ Doubling of serum creatinine to a concentration ≥ 2.5 mg/dL within 2 weeks ■ No response to diuretic withdrawal and 2 day fluid challenge with 1 g/kg/day of albumin 20-25% ■ Cirrhosis with ascites ■ Absence of shock ■ No current or recent use of nephrotoxic drugs (NSAIDs, contrast dye, etc) ■ No signs of structural kidney injury – Absence of proteinuria (>500 mg/day) – Absence of hematuria (>50 RBCs per high power field) – Normal findings on renal ultrasonography	HRS-AKI ■ Increase in serum creatinine of ≥ 0.3 mg/dL within 48 hours - OR ■ Increase in serum creatinine ≥ 1.5 times from baseline (creatinine value within previous 3 months, when available, may be used as baseline, and value closest to presentation should be used) ■ No response to diuretic withdrawal and 2 day fluid challenge with 1 g/kg/day of albumin 20-25% ■ Cirrhosis with ascites ■ Absence of shock ■ No current or recent use of nephrotoxic drugs (NSAIDs, contrast dye, etc) ■ No signs of structural kidney injury – Absence of proteinuria (>500 mg/day) – Absence of hematuria (>50 RBCs per high power field) – Normal findings on renal ultrasound
HRS type 2 ■ Gradual increase in serum creatinine, not meeting criteria above	HRS-NAKI HRS-AKD ■ Estimated glomerular filtration rate <60 mL/min/1.73 m² for <3 months in absence of other potential causes of kidney disease ■ Percentage increase in serum creatinine $<50\%$ using last available value of outpatient serum creatinine within 3 months as baseline value HRS-CKD ■ Estimated glomerular filtration rate <60 mL/min/1.73 m² for ≥ 3 months in absence of other potential causes of kidney disease

Fig 1 | Previous and current definitions of hepatorenal syndrome (HRS). AKD=acute kidney disease; AKI=acute kidney injury; CKD=chronic kidney disease; NAKI=non-acute kidney injury; NSAID=non-steroidal anti-inflammatory drug; RBC=red blood cell

increased production of vasodilators, including nitric oxide, carbon monoxide, prostacyclins, and endocannabinoids, in the splanchnic circulation. Systemic vasodilation results in reduction of effective arterial blood volume (EABV) and systemic arterial pressure. With progression of liver disease, reduction in cardiac output often precedes the development of hepatorenal syndrome, while the peripheral vascular resistance remains unchanged.²⁶ These findings suggest a role for cirrhotic cardiomyopathy in the pathogenesis of hepatorenal syndrome.

Systemic vasoconstrictor pathways, such as the renin-angiotensin-aldosterone system, sympathetic nervous system, and arginine vasopressin, are activated as a means of increasing the EABV. These mechanisms result in sodium retention, impaired solute-free water excretion and renal vasoconstriction, and, consequently, reduced renal blood flow (fig 2).

In earlier stages, the kidneys are able to maintain a normal glomerular filtration rate owing to

the vasodilatory effect of renal prostaglandins (prostaglandin I2 and prostaglandin E2) on afferent renal arterioles. Thus, glomerular pressure can initially be maintained despite reduced renal blood flow. This balance is disrupted with progression of liver disease and by drugs such as non-steroidal anti-inflammatory agents that inhibit prostaglandin synthesis and reduce filtration fraction, leading to AKI.

Elevated ammonia in cirrhosis disrupts metabolism of arginine, an essential amino acid for the synthesis of nitric oxide.²⁷ Decreased availability of nitric oxide in the renal microcirculation contributes to impaired renal blood flow and consequent functional and ischemic kidney injury.

Systemic inflammation

Systemic inflammatory response syndrome has been observed in almost half of patients with AKI-HRS, independent of the presence of infection.²⁸ Moreover, plasma concentrations of pro-inflammatory




Fig 2 | Pathophysiology of hepatorenal syndrome

cytokines (interleukin-6, tumor necrosis factor- α (TNF- α), vascular cell adhesion protein-1, and interleukin-8) and urinary concentrations of monocyte chemoattractant protein-1 are increased in patients with AKI-HRS compared with those with decompensated cirrhosis without AKI and patients with AKI secondary to pre-renal azotemia.²⁹

Inflammation in cirrhosis is driven by two groups of molecules: pathogen associated molecular patterns (PAMPs) and damage associated molecular patterns (DAMPs). PAMPs represent bacterial products, such as lipopolysaccharide, flagellin, and nigericin, which result from translocation of gut bacteria or bacterial infections, whereas DAMPs represent intracellular components released from injured hepatocytes, including high mobility group protein B1, heat shock protein, adenosine triphosphate, double stranded genomic DNA, and others. In the absence of overt bacterial infection, both PAMPs and DAMPs may drive inflammation and release of pro-inflammatory cytokines through activation of pattern recognition receptors such as toll-like receptors (TLRs). The systemic pro-inflammatory response, in turn, may lead to increased arterial production of vasodilators (such as nitric oxide) and, consequently, further reduction in systemic vascular resistance and EABV.

In addition to a systemic effect, DAMPs and PAMPs may have a direct role in the kidneys. Patients with cirrhosis and renal dysfunction show increased expression of TLR4 receptors and caspase-3 in tubular renal cells, both important components of

the innate immune system.³⁰ Gut decontamination has been shown to reduce renal expression of TLR4 and prevent renal dysfunction and tubular damage in animal models of cirrhosis, suggesting that increased TLR4 expression in the kidneys may result from exposure to PAMPs.³¹ Expression of TLR4 and other pattern recognition receptors may also increase as a result of renal ischemia secondary to reduced renal blood flow in hepatorenal syndrome.^{32–33} In experimental models of ischemic AKI, innate immunity has been shown to be responsible for the renal damage.³⁴ Nevertheless, further studies are needed to elucidate the exact mechanisms by which systemic inflammation leads to hepatorenal syndrome.

Hepato-adrenal syndrome

Relative adrenal insufficiency (RAI) is present in 24–47% of patients with decompensated cirrhosis and ascites and may play a role in the development of hepatorenal syndrome.^{35–37} Compared with patients with normal adrenal function, those with RAI have lower arterial pressure and higher serum concentrations of renin and noradrenaline and are at increased risk of AKI-HRS, sepsis, and short term mortality.^{37–40}

The mechanisms leading to the development of hepato-adrenal syndrome remain elusive but may relate to exhaustion of substrates for synthesis of cortisol and impairment of the hypothalamus-pituitary axis by circulating PAMPs and pro-

inflammatory cytokines.^{41 42} Treatment with hydrocortisone may improve outcomes in patients with RAI and septic shock, including shock resolution and short term survival.⁴³ However, the effect of glucocorticoid replacement therapy for hepatorenal syndrome in the absence of shock and its role in prevention and treatment of hepatorenal syndrome remain to be determined.

Cholemic (or bile cast) nephropathy

Cholemic nephropathy has long been reported in patients with cirrhosis and high serum bilirubin concentrations and is thought to be caused by formation of obstructing intratubular bile acid casts and direct bile acid toxicity to tubular cells. Histopathologic studies have shown the presence of intratubular bile acid casts in 18-75% of patients with AKI-HRS.^{44 45} Urinary bilirubin and urobilinogen are elevated in most patients and may be clues to the diagnosis.⁴⁴ Cholemic nephropathy may be present in most patients with jaundice and AKI-HRS and possibly affects outcomes and treatment response, but the prevalence is likely underestimated. Serum bilirubin concentrations above 10 mg/dL have been associated with a lower rate of response to vasoconstrictors in patients with AKI-HRS compared with patients with serum bilirubin below 10 mg/dL (13% v 67%; $P=0.001$).^{46 47} Treatments targeting bile acids in patients with jaundice and AKI-HRS may therefore be beneficial. A recent study in an experimental model of biliary cirrhosis showed that administration of norursodeoxycholic acid ameliorates renal function and histologic findings and may represent a therapeutic option for cholemic nephropathy.⁴⁸ Nevertheless, the pathophysiologic contribution of cholemic nephropathy to AKI-HRS has not been convincingly demonstrated and needs further investigation.

Intra-abdominal hypertension

Elevated intra-abdominal pressure (>12 mm Hg) is an underappreciated cause of AKI and may play a role in the development of hepatorenal syndrome in patients with refractory ascites. A small study assessing the short term effects of paracentesis in patients with hepatorenal syndrome showed a significant improvement in creatinine clearance after reduction of intra-abdominal pressure.⁴⁹ However, careful monitoring of systemic hemodynamic parameters with guided plasma expansion to prevent development of post-paracentesis circulatory dysfunction is paramount. The use of bedside echocardiography for estimation of inferior vena cava diameter and collapsibility may help to determine whether intra-abdominal hypertension might be contributing to renal dysfunction and, thus, help to identify which patients may benefit from large volume paracenteses.⁵⁰

Hepatorenal reflex hypothesis

A link between the kidneys and the liver has long been proposed, suggesting the presence of

osmoreceptors, chemoreceptors, and baroreceptors in the liver that directly affect kidney function through complex neural circuits.⁵¹ This hypothesis has been supported by several experimental studies through manipulation of portal blood osmolality and/or chemical composition, as well as changes in portal pressure. One animal study showed that activation of intrahepatic adenosine receptors in response to reduced portal venous blood flow results in renal sodium and water retention, similar to that observed in hepatorenal syndrome.⁵² Although these findings may have important clinical implications, further studies translating these results to humans are needed.

Differential diagnosis and biomarkers

One important diagnostic criterion for AKI-HRS is exclusion of structural kidney injury, which relies on urine microscopy and urine sodium excretion. Pre-renal azotemia represents the leading cause of AKI in patients with cirrhosis (46-66% of all cases), owing to frequent use of diuretics, large volume paracenteses without albumin, gastrointestinal bleeding, and gastrointestinal fluid losses secondary to lactulose induced diarrhea.^{53 54} The reported prevalence of AKI-HRS and acute tubular necrosis (ATN) varies widely, likely reflecting the challenge in differentiating the two conditions. The diagnosis of AKI-HRS requires the absence of shock, proteinuria (>500 mg/day), and microhematuria (>50 red blood cells per high power field), along with normal renal ultrasonography. However, patients who fulfill these criteria may still have tubular damage, so ATN cannot be confidently excluded.

Some experts have used urinary sodium (>40 mEq/L), fractional excretion of sodium (FeNa >2%), and low urine osmolality (<400 mOsm/L) as suggestive of ATN; however, urinary sodium can be elevated secondary to diuretics, frequently used in this group of patients with large volume ascites.⁵⁵ Conversely, low FeNa has also been observed in patients with biopsy proven ATN,⁵⁶ so urinary sodium and FeNa are no longer part of the diagnostic criteria of AKI-HRS. Studies in patients without liver disease have found that fractional excretion of urea is superior to FeNa in differentiating AKI-HRS from ATN in patients taking diuretics.⁵⁷ Reabsorption of urea, as opposed to sodium, occurs primarily in the proximal renal tubules and, therefore, is not affected by diuretics such as furosemide and spironolactone, which act in the loop of Henle and distal tubules, respectively.⁵⁷ In a recent study assessing the role of fractional excretion of urea in derivation and validation cohorts of patients with cirrhosis and AKI taking diuretics, the area under the receiver operating characteristic curve to differentiate ATN from other causes (cut-off 33.4, sensitivity 85, and specificity 100) was 0.96.⁵⁸

Novel urine biomarkers of tubular injury have long been sought to differentiate AKI-HRS and ATN in patients with cirrhosis. Candidate biomarkers include tubular proteins released during cell damage

(N-acetyl- β -D-glucosaminidase, α -glutathione S-transferase), tubular proteins up regulated by injury (kidney injury molecule-1, neutrophil gelatinase associated lipocalin (NGAL), liver-type fatty acid binding protein), plasma proteins with diminished tubular reabsorption (α -1-microglobulin, β -2-microglobulin, retinol binding protein), and markers of inflammation (interleukin-18).^{59–60} Among these, NGAL has been the most widely studied biomarker in patients with cirrhosis and has shown the greatest diagnostic accuracy in differentiating ATN from AKI-HRS.^{61–66} Urinary NGAL seems to be superior to plasma concentrations and performs better when measured after the two day volume challenge recommended in the management of AKI. The urinary NGAL cut-off value of 220 μ g/g of creatinine obtained after the fluid challenge has the highest diagnostic accuracy for ATN (odds ratio 42.9, 95% confidence interval 13.9 to 132.3).⁶⁴ Moreover, urinary NGAL is an independent predictor of short term mortality.^{62–64–66} One of the limitations of biomarker studies in this population is the lack of a gold standard short of kidney biopsy. Because of the inherent risks, kidney biopsies are not routinely obtained in clinical practice, and human studies rely on expert adjudication for differentiating ATN from AKI-HRS. Nevertheless, the results are encouraging and justify further study.

Risk factors and prevention

Hyponatremia, high plasma renin activity, and liver size,⁶⁷ as well as severity of ascites,⁶⁸ are predictors of AKI-HRS. Acute hemodynamic changes associated with infections and large volume paracentesis without albumin administration represent the most common precipitants of AKI-HRS. The prevalence of unprecipitated AKI is low (1.8%).⁶⁸ AKI-HRS develops in as many as 30% of patients with spontaneous bacterial peritonitis (SBP),⁶⁹ as well as other infections,^{70–71} and is associated with worse outcomes.^{47–72} Infection associated AKI-HRS may be prevented by administration of intravenous albumin in addition to antibiotic treatment in the setting of SBP (8.3% v 30.6% with antibiotics alone; $P=0.01$) and may reduce overall mortality (16% v 35.4%; odds ratio 0.34, 0.19 to 0.60).^{73–74} Albumin administration in patients with infections other than SBP may also improve circulatory function and delay the development of renal dysfunction,⁷⁵ but it has not been shown to prevent AKI-HRS or improve survival.^{76–78}

Long term use of weekly albumin in patients with decompensated cirrhosis and ascites has been assessed in a large RCT ($n=431$). Compared with standard of care, the addition of weekly albumin for 18 months improved overall survival (77% v 66%; $P=0.028$) and also reduced the incidence of hepatorenal syndrome (odds ratio 0.39, 0.19 to 0.76).⁷⁹ In contrast, a similar trial evaluating the long term use of albumin and midodrine in 196 patients with decompensated cirrhosis on the waitlist for liver transplantation failed to show a survival benefit at

one year, and complications of cirrhosis were not prevented. However, in this latter study only a small number of patients completed a full year on the study and the median length of treatment was only 80 days.⁸⁰ The forthcoming PRECIOUSA12 (Effects of Long term Administration of Human Albumin in Subjects With Decompensated Cirrhosis and Ascites) trial will hopefully elucidate the role of long term albumin use in this population.

Antibiotic prophylaxis in patients at risk of SBP, as determined by low ascitic fluid protein (<1.5 mg/dL) associated with liver and/or kidney dysfunction (bilirubin >3 mg/dL, serum sodium <130 mEq/L, Child-Turcotte-Pugh score >10 , and/or serum creatinine >1.2 mg/dL), not only prevents development of SBP but also significantly reduces the risk of AKI-HRS and overall mortality.^{81–82}

Post-paracentesis circulatory dysfunction occurs after large volume paracenteses (≥ 4.5 L) and is associated with hypotension, hyponatremia, and increased risk of AKI-HRS.⁸³ Albumin administration following a large volume paracentesis significantly reduces this risk and improves overall survival in these patients.^{84–85} This protective effect seems to be unique to albumin, compared with other volume expanders,⁸⁵ which suggests an additional benefit of albumin beyond simply volume expansion.⁸⁶ Albumin has important antioxidant and anti-inflammatory properties and helps to stabilize endothelial function.^{87–88} Moreover, human albumin is able to bind and potentially inactivate endotoxins (lipopolysaccharides) and so may reduce their negative effect on circulatory and kidney functions.⁸⁹ Given its potential immunomodulatory effects,⁹⁰ albumin is being investigated in an RCT for prevention of infection in cirrhosis.⁹¹

Management

The updated diagnostic criteria, with removal of a minimum serum creatinine concentration, allow for earlier diagnosis and treatment of AKI-HRS. Rather than waiting for a doubling of creatinine to reach 2.5 g/dL, drug treatment may now be started immediately after an unsuccessful fluid challenge. This is likely to result in higher reversal rates and better outcomes, as response to vasoconstrictors is dependent on the serum creatinine concentration at the start of treatment.²² Nevertheless, AKI stage 1A (serum creatinine <1.5 g/dL) is most often secondary to hypovolemia and is expected to resolve in more than 90% of patients at this stage, compared with a half of patients with stage 1B disease (serum creatinine ≥ 1.5 g/dL).^{3–54} Therefore, European Association for the Study of the Liver (EASL) guidelines recommend reserving the use of vasoconstrictors for patients with AKI-HRS stage 1B or greater (fig 3).⁹² However, in most countries, the use of vasoconstrictors is indicated for hepatorenal syndrome type 1, based on the old definition, and the use of vasoconstrictors in patients with creatinine below 2.5 mg/dL is considered off-label.

Once a diagnosis of AKI is made, management of hepatorenal syndrome starts with a fluid challenge

of 20-25% intravenous albumin at 1 g/kg/day for two days and withdrawal of diuretics. This is not only needed to rule out pre-renal azotemia but also promotes early plasma volume expansion in the setting of reduced EABV. This initial phase also includes temporary discontinuation of non-selective β blockers given their negative inotropic effect, which reduces cardiac output.^{93 94} These should be carefully reinstituted once renal function and mean arterial pressure (MAP) improve.⁹³

The specific treatment of AKI-HRS comprises vasoconstrictors in combination with albumin infusion and reversal of precipitant factors. Bacterial infections, particularly SBP, should be ruled out by blood, urine, and ascitic fluid cultures and a chest radiograph. Although antibiotics may help to prevent development of AKI-HRS,^{81 95 96} their benefit in patients with established AKI-HRS in the absence of infection has not been demonstrated.

Vasoconstrictors

Splanchnic vasoconstriction in patients with cirrhosis results in a reduction in portal pressure and an increase in EABV and renal blood flow, especially when combined with intravenous administration of albumin. Renal perfusion directly correlates

with changes in MAP and is negatively affected by intra-abdominal pressure driven by ascites. A significant increase in MAP promoted by the use of vasoconstrictors is associated with a higher likelihood of reversal of hepatorenal syndrome.⁹⁷

Several RCTs have confirmed the efficacy of vasoconstrictors, which represent the mainstay treatment of AKI-HRS.⁹⁸ The available options include terlipressin, noradrenaline, and the combination of midodrine plus octreotide.

Terlipressin is not yet available in North America but can be prescribed to treat AKI-HRS in many European and Asian countries. Terlipressin, a synthetic vasopressin analog with predominant vasopressin 1A receptor effect, acts primarily as a splanchnic vasoconstrictor.⁹⁹ In addition, terlipressin activates vasopressin 1B receptors, which stimulate release of adrenocorticotrophic hormone and cortisol and might counteract the relative adrenal insufficiency commonly observed in patients with decompensated cirrhosis.¹⁰⁰ Furthermore, terlipressin shows greater efficacy in reversal of AKI-HRS in patients with a systemic inflammatory response,¹⁰¹ which may relate to indirect vasopressin mediated anti-inflammatory effects.¹⁰² An agent with selective vasopressin 1 activity is preferable in patients with cirrhosis to




Fig 3 | Algorithm for management of acute kidney injury (AKI) in patients with cirrhosis. ATN=acute tubular necrosis; HRS=hepatorenal syndrome.

*Risk factor management: hold nephrotoxic drugs/ β blockers, withdraw diuretics, treat infections, plasma volume expansion as needed

prevent unwanted solute-free water absorption and consequent worsening hyponatremia and volume overload induced by renal vasopressin 2 receptor activation.¹⁰³ Because terlipressin is also a modest vasopressin 2 receptor agonist,⁹⁹ an acute reduction in serum sodium concentration may occur, although this effect is mainly observed in patients with cirrhosis without ascites treated with terlipressin for variceal bleeding.¹⁰⁴ A novel selective vasopressin 1A receptor agonist, selepressin, has recently been investigated as an alternative to vasopressin in management of septic shock and has shown a lower risk of hyponatremia, volume overload, and pulmonary edema in animal models of sepsis.^{105 106} The role of selepressin in hepatorenal syndrome warrants further study.

Norepinephrine (intravenous) and midodrine (oral) are systemic vasoconstrictors through activation of α -1 adrenergic receptors on vascular smooth muscle cells. Octreotide, a somatostatin analog, acts by inhibiting secretion of glucagon, a splanchnic vasodilator, and is a direct mesenteric vasoconstrictor.¹⁰⁷ However, the effect of octreotide in patients with cirrhosis is dampened by antagonistic effects of local nitric oxide release,¹⁰⁸ and octreotide alone has limited benefit in hepatorenal syndrome.¹⁰⁹ Midodrine monotherapy also does not improve renal function in patients with hepatorenal syndrome despite its hemodynamic effects.¹¹⁰ In contrast, a combination of octreotide and midodrine has potential benefit in hepatorenal syndrome and has become the standard of care in countries where terlipressin is not yet approved.^{111 112}

Efficacy

Comparative studies (some of them open label) evaluating terlipressin, norepinephrine, and/or octreotide/midodrine have found terlipressin, in combination with intravenous albumin, to be the most effective drug treatment for AKI-HRS (table 1).^{23 113-122} The efficacy of terlipressin plus albumin in achieving complete reversal of hepatorenal syndrome, defined as at least 50% reduction in serum creatinine to a final value below 1.5 mg/dL, ranges from 19% to 56% compared with 3-14% with albumin alone.^{23 101 115 118 121} Conversely, administration of terlipressin alone is markedly inferior to a combination of terlipressin and albumin (complete hepatorenal syndrome reversal rates of 25% and 77%, respectively).¹²³ Two large European trials and one North American trial have shown efficacy for terlipressin.^{23 115 124} However, in the REVERSE study, a large multicenter phase III trial, terlipressin led to complete reversal of hepatorenal syndrome in only 19.6% of patients compared with 13.1% in the placebo group ($P=0.22$).¹¹⁶ One of the potential reasons for the discrepant results was the requirement for a confirmatory creatinine concentration below 1.5 mg/dL 48 hours after initial hepatorenal syndrome reversal, which was not available in many patients owing to discharge from hospital or transplantation. Another reason could

be the short duration of treatment with terlipressin in a large number of patients. This has led to the execution of yet another RCT in North America designed to overcome the methodological obstacles encountered in the REVERSE trial. The CONFIRM study has been published only in abstract form. The verified hepatorenal syndrome reversal rate with terlipressin plus albumin was 29.1% compared with 15.8% with albumin plus placebo ($P<0.012$).¹¹⁷

Norepinephrine is also effective and safe in AKI-HRS,¹²⁵ with similar rates of hepatorenal syndrome reversal to terlipressin in small randomized studies, ranging between 39% and 70%.¹¹⁸⁻¹²⁰ Norepinephrine is less expensive than terlipressin; however, central line placement and admission to an intensive care unit are needed for administration, which may offset the cost benefit. Terlipressin may be given through a peripheral line on the medical floor.¹¹⁹

Only one study has directly compared combination midodrine/octreotide with terlipressin.¹²¹ The complete response rate for midodrine/octreotide was only 4.8% compared with 55.5% with terlipressin, and the overall response (complete or partial) rates were 28.6% and 70.4%, respectively. Although no placebo controlled studies have been done with either norepinephrine or midodrine/octreotide, a recent network meta-analysis has allowed for indirect comparisons across trials. Compared with placebo, combination midodrine/octreotide was not significantly superior in achieving reversal of hepatorenal syndrome (odds ratio 0.44, 0.06 to 3.23), whereas norepinephrine and terlipressin were both superior to midodrine/octreotide and placebo and were equally effective.⁹⁸

Side effects

The most common side effects of terlipressin are diarrhea and abdominal pain, reported in 10-20% of patients overall. Discontinuation of terlipressin due to serious adverse events is needed in 4-22% (median 8%) of patients, with a rate of peripheral ischemic events between 4% and 13%. The rate of myocardial infarction or intestinal ischemia ranges from 2% to 13%.⁹⁸ Pulmonary edema may occur in patients with hepatorenal syndrome treated with terlipressin and albumin, and volume status should be carefully monitored in these patients.¹¹⁶ Continuous terlipressin infusion is associated with a lower rate of adverse events compared with bolus administration, likely owing to lower dosing needed.¹²⁴ Nevertheless, patients should be monitored at least twice daily for signs of ischemia in skin, tongue, and fingers while on therapy, and terlipressin should be avoided in patients with a history of coronary artery disease or peripheral vascular disease.⁹² Discontinuation of treatment due to adverse events is less common with norepinephrine and midodrine/octreotide; however, tachyarrhythmias or bradycardia can be seen (table 2).¹²¹

Predictors of treatment response

Several factors have been shown to negatively affect response of hepatorenal syndrome to drug

Table 1 | Study design and outcomes of randomized controlled trials of vasoactive drugs for treatment of hepatorenal syndrome-acute kidney injury (HRS-AKI)

Study	Trial design	Drug comparisons (No of patients)	No (%) HRS reversal	No (%) mortality
Terlipressin versus placebo/control				
Solanki et al, 2003 ¹¹³	Single center, single blind, placebo controlled	Terlipressin 1 mg every 12 h for 15 days (n=12) v placebo (n=12)	NA	Terlipressin 7/12 (58.3) v placebo 12/12 (100)
Neri et al, 2008 ¹¹⁴	Single center, open label	Terlipressin 1 mg every 8 h for 5 days followed by 0.5 mg every 8 h for 14 days (n=26) v albumin only for 15 days (n=26)	Terlipressin 21/26 (80) v control 5/26 (19)	Terlipressin 7/26 (26.9) v control: 15/26 (57.7)
Sanyal et al, 2008 ¹¹⁵	Multicenter, double blind, placebo controlled	Terlipressin 1 mg every 6 h up to 2 mg every 6 hours for 14 days (n=56) v placebo for 14 days (n=56)	Terlipressin 19/56 (33.9) v placebo 7/56 (12.5)	Terlipressin 32/56 (57.1) v placebo 35/56 (62.5)
Martin-Llahi et al, 2008 ²³	Multicenter, open label	Terlipressin 1 mg every 4 h up to 2 mg every 4 h for 15 days (n=17) v albumin only daily for 15 days (n=18)	Terlipressin 6/17 (35.3) v control 2/18 (11.1)	Terlipressin 17/23 (73.9) v control: 19/23 (82.6)
Boyer et al, 2016 ¹¹⁶	Multicenter, double blind, placebo controlled	Terlipressin 1 mg every 6 h up to 2 mg every 6 h for 14 days (n=97) v placebo for 14 days (n=99)	Terlipressin 19/97 (19.6) v placebo 13/99 (13.1)	Terlipressin 32/97 (33) v placebo: 35/99 (35.3)
Wong et al, 2019 ^{*117}	Multicenter, double blind, placebo controlled	Terlipressin 1 mg every 6 h until verified HRS reversal or for maximum 14 days (n=199) v placebo until verified HRS reversal or for maximum 14 days (n=101)	Terlipressin 58/199 (29.1) v placebo 16/101 (15.8)	Terlipressin 145/199 (72.9) v placebo 72/101 (71.3)
Terlipressin versus norepinephrine				
Alessandria et al, 2007 ¹¹⁸	Single center, open label	Terlipressin 1 mg every 4 h up to 2 mg every 4 h until HRS reversal or for maximum 14 days (n=4) v norepinephrine 0.1 µg/kg/min up to 0.7 µg/kg/min until HRS reversal or maximum 14 days (n=5)	Terlipressin 3/4 (75) v norepinephrine 4/5 (80)	Terlipressin 1/4 (25) v norepinephrine 1/5 (20)
Sharma et al, 2008 ¹¹⁹	Single center, open label	Terlipressin 0.5 mg every 6 h up to 2 mg every 6 h for 15 days (n=20) v norepinephrine 0.5 mg/h up to 3 mg/h for 15 days (n=20)	Terlipressin 8/20 (40) v norepinephrine 10/20 (50)	Terlipressin 9/20 (45) v norepinephrine 9/20 (45)
Singh et al, 2012 ¹²⁰	Single center, open label	Terlipressin 0.5 mg every 6 h up to 2 mg every 6 h until HRS reversal or for maximum 14 days (n=23) v norepinephrine 0.5 mg/h up to 3 mg/h until HRS reversal or for maximum 14 days (n=23)	Terlipressin 9/23 (39.1) v norepinephrine 10/23 (43.5)	Terlipressin 16/23 (69.5) v norepinephrine 15/23 (65.2)
Terlipressin versus midodrine plus octreotide				
Cavallin et al, 2015 ¹²¹	Multicenter, open label	Terlipressin 3-12 mg per 24 h until HRS reversal or for maximum 14 days (n=27) v midodrine 7.5-12.5 mg every 8 h orally plus octreotide 100-200 µg every 8 h subcutaneously until HRS reversal or for maximum of 14 days (n=22)	Terlipressin 15/27 (55.5) v midodrine plus octreotide 1/22 (4.5)	Terlipressin 8/27 (29.6) v midodrine plus octreotide 7/22 (31.8)

NA=not available.

*Data published in abstract format.

treatment, including model for end stage liver disease (MELD) score,¹²⁶ pretreatment serum creatinine concentration,^{22 23} sepsis,¹²⁶ extrahepatic organ failure,¹²⁷ and systemic inflammation.²⁹ Lower serum creatinine concentrations at the start of treatment are associated with higher rates of HRS reversal, whereas only a negligible response is expected when creatinine exceeds 7 g/dL.^{22 23} Thus, earlier diagnosis of AKI-HRS with the elimination of creatinine concentration cut-off, will likely result in higher rates of response to treatment. Total serum bilirubin concentration has also been found to predict response, with an area under the receiver operating characteristic curve of 0.77 (P<0.0001) for a bilirubin cut-off of 10 mg/dL.⁴⁶ These findings suggest the presence of structural damage.

Finally, response to treatment with vasoactive agents directly correlates with sustained increase in MAP.⁹⁷ Independent of the agents used, improvement in renal function is preceded by a sustained rise in MAP by 5-10 mm Hg on average from baseline.^{22 46 111}

Albumin

Albumin infusion is essential for effective management of AKI-HRS. Several studies provide supportive evidence for a multifaceted mode of action by albumin, which may also include volume expansion and positive cardiac inotropic effect,¹²⁸

as well as antioxidant and immunomodulatory properties.^{88 129 130} In a single non-randomized study comparing terlipressin plus albumin versus terlipressin alone, the combination with albumin resulted in a significantly higher response (77% v 25%; P=0.03).¹²³ In addition to plasma volume expansion and consequent increase in EABV, albumin has shown several other benefits in hepatorenal syndrome. In a study comparing hydroxyethyl starch, a synthetic colloid, with albumin in patients with SBP, albumin use was associated with lower plasma concentrations of von Willebrand related antigen and factor VIII, suggesting that albumin, but not hydroxyethyl starch, may reduce endothelial activation.⁸⁶

One important property of albumin is its ability to bind a wide range of substances including bile acids, hormones, cytokines, long chain fatty acids, nitric oxide, endotoxin, and other bacterial products.¹³¹ This is the basis for the advent of molecular absorbent recirculatory systems, a modified dialysis method that clears substances bound to albumin. This results in a significant reduction in serum creatinine concentrations in patients with hepatorenal syndrome.¹³² However, improvement in renal function and systemic hemodynamics is not observed in patients with hepatorenal syndrome refractory to vasoconstrictors, despite reduction in nitric oxide concentrations.¹³³

Table 2 | Vasoconstrictors in hepatorenal syndrome-acute kidney injury

Treatment	Dose	Route	Frequency	Side effects
Midodrine and octreotide	7.5-15 mg and 100-200 µg	Oral and subcutaneous	Three times daily	Bradycardias, paresthesias, abdominal pain, diarrhea, cholelithiasis, hyperglycemia
Norepinephrine	0.5-3 mg/h; titrate to achieve 10 mm Hg increase in mean arterial pressure	Intravenous	Continuous infusion	Nausea, vomiting, anxiety, cardiac dysrhythmias
Terlipressin	1 mg; titrate if no improvement (decrease in serum creatinine by 25% by day 3) up to maximum 12 mg/day	Intravenous	Every 4-6 h or continuous infusion	Diarrhea, abdominal pain, peripheral ischemia, myocardial infarction, mesenteric ischemia, pulmonary edema

Transjugular intrahepatic portosystemic shunt

The creation of an intrahepatic shunt aimed at reducing portal pressure has shown significant benefit in patients with cirrhosis who cannot tolerate diuretics or have refractory ascites and who have uncontrolled variceal bleeding.^{134 135} However, only a few studies have explored the role of transjugular intrahepatic portosystemic shunt (TIPS) insertion in AKI-HRS, so its use remains investigational in this group. One small non-randomized study showed significant improvement in renal function after TIPS insertion for AKI-HRS, with reduction in plasma

renin activity and aldosterone and norepinephrine concentrations.¹³⁶ A meta-analysis including nine studies in which 128 patients with hepatorenal syndrome underwent TIPS insertion (77 patients with hepatorenal syndrome type 1 and 51 patients with type 2), showed significant improvement in serum creatinine, serum sodium, and urine output.¹³⁷ Patients with markedly elevated bilirubin, active infection, or overt hepatic encephalopathy were excluded, so the role of TIPS in hepatorenal syndrome may be limited to a highly selected group of patients. On the other hand, TIPS may have a greater

Table 3 | Recommendations adapted from American Association for the Study of Liver Disease and European Association for the Study of the Liver^{92 149}

Recommendation	Strength of recommendation
American Association for the Study of Liver Diseases guidelines 2012	
Urinary biomarkers such as neutrophil gelatinase associated lipocalin may assist in the differential diagnosis of azotemia in patients with cirrhosis	Weak
Albumin infusion plus administration of vasoactive drugs such as octreotide and midodrine should be considered in the treatment of type I hepatorenal syndrome	Weak
Albumin infusion plus administration of norepinephrine should also be considered in the treatment of type I hepatorenal syndrome, when the patient is in the intensive care unit	Weak
Patients with cirrhosis, ascites, and type I or type II hepatorenal syndrome should have an expedited referral for liver transplantation	Strong
European Association for the Study of the Liver guidelines 2018	
Vasoconstrictors and albumin are recommended in all patients meeting the current definition of AKI-HRS stage >1A. Such patients should be expeditiously treated with vasoconstrictors and albumin	Strong
Terlipressin plus albumin should be considered as the first line therapeutic option for the treatment of HRS-AKI. Terlipressin can be used by iv boluses at the initial dose of 1 mg every 4-6 h. However, giving terlipressin by continuous iv infusion at initial dose of 2 mg/day makes reduction of the global daily dose of the drug, and thus the rate of its adverse effects, possible. In case of non-response (decrease in SCr <25% from the peak value), after two days, the dose of terlipressin should be increased in a stepwise manner to a maximum of 12 mg/day	Strong
Albumin solution (20%) should be used at a dose of 20-40 g/day. Ideally, apart from routinely monitoring patients with HRS-AKI, the serial measurement of CVP or other measures of assessing central blood volume can help to prevent circulatory overload by optimizing the fluid balance and helping to titrate the dose of albumin	Strong
Noradrenaline can be an alternative to terlipressin. However, limited information is available	Weak
In contrast to terlipressin, the use of noradrenaline always requires a central venous line and, in several countries, the transfer of the patient to an ICU. Midodrine plus octreotide can be an option only when terlipressin and noradrenaline are unavailable, but its efficacy is much lower than that of terlipressin	Strong
According to the new definition of HRS-AKI, complete response to treatment should be defined by a final SCr within 0.3 mg/dL (26.5 µmol/L) from the baseline value, while partial response should be defined by the regression of AKI stage to a final SCr ≥0.3 mg/dL (26.5 µmol/L) from the baseline value	Strong
Adverse events related to terlipressin or noradrenaline include ischemic and cardiovascular events. Thus, a careful clinical screening including electrocardiography is recommended before starting the treatment. Patients can be treated on a regular ward, but the decision to transfer to higher dependency care should be case based. For the duration of treatment, close monitoring of patients is important. According to the type and severity of side effects, treatment should be modified or discontinued	Strong
In cases of recurrence of HRS-AKI on cessation of treatment, a repeat course of therapy should be given	Strong
Terlipressin plus albumin is also effective in the treatment of HRS outside the criteria of AKI (HRS-NAKI), formerly known as HRS type II. Unfortunately, recurrence after the withdrawal of treatment is the norm, and data on the effect of the treatment on long term clinical outcome are controversial, particularly from the perspective of LT. As such, vasoconstrictors and albumin are not recommended in this clinical scenario	Strong
Insufficient data exist to allow TIPS to be advocated in HRS-AKI, but it could be suggested in selected patients with HRS-NAKI	Weak
LT is the best therapeutic option for patients with HRS regardless of the response to drug therapy	Strong
The decision to initiate RRT should be based on the individual severity of illness	Weak
The indication for liver-kidney transplantation remains controversial. This procedure should be considered in patients with significant CKD or with sustained AKI including HRS-AKI with no response to drug therapy	Strong
Albumin (1.5 g/kg at diagnosis and 1 g/kg on day 3) should be given in patients with SBP to prevent AKI	Strong
Norfloxacin (400 mg/day) should be given as prophylaxis of SBP to prevent HRS-AKI	Strong

AKI=acute kidney injury; CKD=chronic kidney disease; CVP=central venous pressure; HRS=hepatorenal syndrome; ICU=intensive care unit; iv=intravenous; LT=liver transplantation; NAKI=non-AKI; RRT=renal replacement therapy; SBP=spontaneous bacterial peritonitis; SCr=serum creatinine; TIPS=transjugular intrahepatic portosystemic shunt.

protective effect as shown by reduced incidence of hepatorenal syndrome in patients with cirrhosis and diuretic refractory ascites post-TIPS.¹³⁴

Renal replacement therapy

Renal replacement therapy (RRT) may be indicated for patients with AKI-HRS unresponsive to drug treatment and with volume overload, uremia, or electrolyte derangements; however, RRT does not improve survival in hepatorenal syndrome,¹³⁸ and it should be reserved for use as a bridge to transplantation, when transplantation is an option.¹³⁹ Short term mortality in patients with cirrhosis and AKI who are ineligible for transplantation approaches 90%,^{140 141} independently of the cause of AKI.¹⁴² Therefore, RRT is often futile in this setting.

Liver transplantation

The functional nature of hepatorenal syndrome means that improvement in renal function is expected with liver transplantation, which remains the optimal treatment of AKI-HRS whenever feasible.¹⁴³⁻¹⁴⁵ However, kidney recovery is not universal and is dependent on multiple factors, particularly duration of kidney injury.¹⁴⁶ In such cases, simultaneous liver-kidney transplantation is recommended rather than liver transplantation alone. However, accurately predicting native kidney recovery after liver transplantation remains a challenge. In the US, the Organ Procurement and Transplantation Network policy for simultaneous liver-kidney organ allocation requires sustained AKI defined as need for dialysis or measured or calculated creatinine clearance or GFR of 25 mL/min or below for a minimum of six consecutive weeks.¹⁴⁷ Despite best efforts, almost 10% of patients with either AKI or CKD who receive a liver alone may have persistent or progressive renal failure after transplant.¹⁴⁸ Patients with ATN are at particularly increased risk of chronic kidney disease (stage 4 or 5) post-transplant, and the lack of ideal biomarkers often results in misdiagnosis.¹⁴⁹ This has led to a consensus “safety net” for prioritization of liver recipients on the kidney waiting list if they are registered within a year after their liver transplant.¹⁵⁰

Guidelines

Prevention, diagnosis, and management of hepatorenal syndrome are included in both the American Association for the Study of Liver Diseases (AASLD) and the EASL guidelines on the management of patients with ascites or decompensated cirrhosis (table 3).^{92 151} The AASLD guidelines were last updated in 2012 and do not contain the most up-to-date diagnostic criteria and classification of hepatorenal syndrome adopted in 2015. In contrast, the European guidelines published in 2018 incorporate the recent changes, including the exclusion of a creatinine cut-off for diagnosis of HRS-AKI. This important change now allows for earlier treatment of HRS-AKI with vasoactive agents

as discussed earlier. Another important difference between the two guidelines is the options for medical management of hepatorenal syndrome. Terlipressin plus albumin is considered first line treatment of HRS-AKI according to the EASL guidelines; however, terlipressin is not yet available in the US, and management of HRS-AKI is limited to albumin, octreotide plus midodrine, or norepinephrine as per the American guidelines.

Emerging treatments

New treatment options are urgently needed, as efficacy of vasoconstrictors and albumin is limited to less than half of patients with AKI-HRS. Serelaxin (recombinant human relaxin-2) is novel agent that acts on renal vasculature and results in increased renal blood flow, reduced renal vascular resistance, and reversal of endothelial dysfunction.¹⁵² Furthermore, serelaxin has been shown to reduce intrahepatic vascular resistance in animal models of cirrhosis, thereby ameliorating portal hypertension.¹⁵³ An exploratory randomized phase II study showed an increase in total renal arterial blood flow by 65% in patients with compensated cirrhosis treated with serelaxin.¹⁵⁴ No studies have been reported in patients with hepatorenal syndrome.

In addition to vasoactive drugs, treatments targeting systemic inflammation, including DAMPs, PAMPs, and downstream signaling, could be explored in hepatorenal syndrome. One small randomized study showed that pentoxifylline, a phosphodiesterase inhibitor with anti-TNF- α activity and anti-inflammatory effect, is safe in patients with AKI-HRS. However, the study failed to show an added benefit compared with standard of care alone (midodrine, octreotide, and albumin).¹⁵⁵ The role of serelaxin and novel therapies targeting inflammation in the clinical management of AKI-HRS needs to be explored in future studies.

Conclusions

Despite advances in biomarker discovery and evolving definitions of hepatorenal syndrome, much of its pathophysiology beyond circulatory dysfunction still remains to be uncovered. Systemic inflammation, in the presence or absence of infection, remains an untapped territory in the understanding of hepatorenal syndrome. Novel translational experimental models of hepatorenal syndrome are needed to fill this gap and, hopefully, will help to identify novel targets for potential drug development. In the meantime, emphasis should be on preventive measures for patients at risk of hepatorenal syndrome, including appropriate antibiotic prophylaxis and albumin use when indicated. For patients with established hepatorenal syndrome, terlipressin with albumin is considered first line medical treatment; however, liver transplantation remains the optimal treatment, and timely referral for transplant evaluation is crucial to avoid permanent kidney damage and the need for simultaneous liver and kidney transplant.

GLOSSARY OF ABBREVIATIONS

- AASLD—American Association for the Study of Liver Diseases
- AKI—acute kidney injury
- AKIN—Acute Kidney Injury Network
- ATN—acute tubular necrosis
- CKD—chronic kidney disease
- DAMP—damage associated molecular pattern
- EABV—effective arterial blood volume
- EASL—European Association for the Study of the Liver
- eGFR—estimated glomerular filtration rate
- FeNa—fractional excretion of sodium
- GFR—glomerular filtration rate
- HRS—hepatorenal syndrome
- ICA—International Club of Ascites
- KDIGO—Kidney Disease Improving Global Outcome
- MAP—mean arterial pressure
- MELD—model for end-stage liver disease
- NAKI—non-acute kidney injury
- NGAL—neutrophil gelatinase associated lipocalin
- PAMP—pathogen associated molecular pattern
- RAI—relative adrenal insufficiency
- RCT—randomized clinical trial
- RIFLE—risk, injury, failure, loss, and end stage kidney disease
- RRT—renal replacement therapy
- SBP—spontaneous bacterial peritonitis
- TIPS—transjugular intrahepatic portosystemic shunt
- TLR—toll-like receptor
- TNF—tumor necrosis factor

QUESTIONS FOR FUTURE RESEARCH

- Can we identify serum and urine biomarkers that allow for earlier diagnosis of hepatorenal syndrome—acute kidney injury (HRS-AKI) and exclusion of acute tubular necrosis?
- What is the optimal stage of HRS-AKI that results in reversal of renal dysfunction and improved survival?
- What is the role of therapies targeting systemic inflammation, including damage associated molecular patterns, pathogen associated molecular patterns, and/or downstream signaling, in the management of hepatorenal syndrome?

Contributors: All authors were involved in the conception, writing and revision of the manuscript.

Competing interests: We have read and understood the BMJ policy on declaration of interests and declare the following interests: PG has received research funding from Mallinckrodt, USA, Grifols, SA, and Gilead, USA, and has worked on advisory boards for Intercept, Gilead, Grifols, Mallinckrodt, and Martin Pharmaceuticals. DAS is a co-investigator of the CONFIRM trial sponsored by Mallinckrodt.

Provenance and peer review: Commissioned; externally peer reviewed.

Patient involvement: No patients were asked for input in the creation of this article.

- 1 Fagundes C, Barreto R, Guevara M, et al. A modified acute kidney injury classification for diagnosis and risk stratification of impairment of kidney function in cirrhosis. *J Hepatol* 2013;59:474-81. doi:10.1016/j.jhep.2013.04.036

- 2 Wu CC, Yeung LK, Tsai WS, et al. Incidence and factors predictive of acute renal failure in patients with advanced liver cirrhosis. *Clin Nephrol* 2006;65:28-33. doi:10.5414/CNP65028
- 3 Piano S, Rosi S, Maresio G, et al. Evaluation of the Acute Kidney Injury Network criteria in hospitalized patients with cirrhosis and ascites. *J Hepatol* 2013;59:482-9. doi:10.1016/j.jhep.2013.03.039
- 4 Sujan R, Cruz-Lemini M, Altamirano J, et al. A Validated Score Predicts Acute Kidney Injury and Survival in Patients With Alcoholic Hepatitis. *Liver Transpl* 2018;24:1655-64. doi:10.1002/lt.25328
- 5 Sherman DS, Fish DN, Teitelbaum I. Assessing renal function in cirrhotic patients: problems and pitfalls. *Am J Kidney Dis* 2003;41:269-78. doi:10.1053/ajkd.2003.50035
- 6 Caregaro L, Menon F, Angeli P, et al. Limitations of serum creatinine level and creatinine clearance as filtration markers in cirrhosis. *Arch Intern Med* 1994;154:201-5. doi:10.1001/archinte.1994.00420020117013
- 7 Mindikoglu AL, Dowling TC, Weir MR, Seliger SL, Christenson RH, Magder LS. Performance of chronic kidney disease epidemiology collaboration creatinine-cystatin C equation for estimating kidney function in cirrhosis. *Hepatology* 2014;59:1532-42. doi:10.1002/hep.26556
- 8 Bellomo R, Ronco C, Kellum JA, Mehta RL, Palevsky P. Acute Dialysis Quality Initiative workgroup. Acute renal failure - definition, outcome measures, animal models, fluid therapy and information technology needs: the Second International Consensus Conference of the Acute Dialysis Quality Initiative (ADQI) Group. *Crit Care* 2004;8:R204-12. doi:10.1186/cc2872
- 9 Mehta RL, Kellum JA, Shah SV, et al. Acute Kidney Injury Network. Acute Kidney Injury Network: report of an initiative to improve outcomes in acute kidney injury. *Crit Care* 2007;11:R31. doi:10.1186/cc5713
- 10 Lassnigg A, Schmidlin D, Mouhieddine M, et al. Minimal changes of serum creatinine predict prognosis in patients after cardiothoracic surgery: a prospective cohort study. *J Am Soc Nephrol* 2004;15:1597-605. doi:10.1097/01.ASN.0000130340.93930.DD
- 11 Chertow GM, Burdick E, Honour M, Bonventre JV, Bates DW. Acute kidney injury, mortality, length of stay, and costs in hospitalized patients. *J Am Soc Nephrol* 2005;16:3365-70. doi:10.1681/ASN.2004090740
- 12 Summary of Recommendation Statements. *Kidney Int Suppl* (2011) 2012;2:8-12. doi:10.1038/kisup.2012.7
- 13 Arroyo V, Ginès P, Gerbes AL, et al. Definition and diagnostic criteria of refractory ascites and hepatorenal syndrome in cirrhosis. International Ascites Club. *Hepatology* 1996;23:164-76. doi:10.1002/hep.510230122
- 14 Belcher JM, Garcia-Tsao G, Sanyal AJ, et al. TRIBE-AKI Consortium. Association of AKI with mortality and complications in hospitalized patients with cirrhosis. *Hepatology* 2013;57:753-62. doi:10.1002/hep.25735
- 15 Tsien CD, Rabie R, Wong F. Acute kidney injury in decompensated cirrhosis. *Gut* 2013;62:131-7. doi:10.1136/gutjnl-2011-301255
- 16 de Carvalho JR, Villela-Nogueira CA, Luiz RR, et al. Acute kidney injury network criteria as a predictor of hospital mortality in cirrhotic patients with ascites. *J Clin Gastroenterol* 2012;46:e21-6. doi:10.1097/MCG.0b013e31822e8e12
- 17 Wong F, O'Leary JG, Reddy KR, et al. North American Consortium for Study of End-Stage Liver Disease. New consensus definition of acute kidney injury accurately predicts 30-day mortality in patients with cirrhosis and infection. *Gastroenterology* 2013;145:1280-8.e1. doi:10.1053/j.gastro.2013.08.051
- 18 Altamirano J, Fagundes C, Dominguez M, et al. Acute kidney injury is an early predictor of mortality for patients with alcoholic hepatitis. *Clin Gastroenterol Hepatol* 2012;10:65-71.e3. doi:10.1016/j.cgh.2011.09.011
- 19 Angeli P, Gines P, Wong F, et al. International Club of Ascites. Diagnosis and management of acute kidney injury in patients with cirrhosis: revised consensus recommendations of the International Club of Ascites. *Gut* 2015;64:531-7. doi:10.1136/gutjnl-2014-308874
- 20 Amathieu R, Al-Khafaji A, Sileanu FE, et al. Significance of oliguria in critically ill patients with chronic liver disease. *Hepatology* 2017;66:1592-600. doi:10.1002/hep.29303
- 21 Angeli P, Garcia-Tsao G, Nadim MK, Parikh CR. News in pathophysiology, definition and classification of hepatorenal syndrome: A step beyond the International Club of Ascites (ICA) consensus document. *J Hepatol* 2019;71:811-22. doi:10.1016/j.jhep.2019.07.002
- 22 Boyer TD, Sanyal AJ, Garcia-Tsao G, et al. Terlipressin Study Group. Predictors of response to terlipressin plus albumin in hepatorenal syndrome (HRS) type 1: relationship of serum creatinine to hemodynamics. *J Hepatol* 2011;55:315-21. doi:10.1016/j.jhep.2010.11.020
- 23 Martín-Llahí M, Pépin MN, Guevara M, et al. TAHRIS Investigators. Terlipressin and albumin vs albumin in patients

- with cirrhosis and hepatorenal syndrome: a randomized study. *Gastroenterology* 2008;134:1352-9. doi:10.1053/j.gastro.2008.02.024
- 24 National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis* 2002;39(Suppl 1):S1-266.
 - 25 Schrier RW, Arroyo V, Bernardi M, Epstein M, Henriksen JH, Rodés J. Peripheral arterial vasodilation hypothesis: a proposal for the initiation of renal sodium and water retention in cirrhosis. *Hepatology* 1988;8:1151-7. doi:10.1002/hep.1840080532
 - 26 Ruiz-del-Arbol L, Monescillo A, Arocena C, et al. Circulatory function and hepatorenal syndrome in cirrhosis. *Hepatology* 2005;42:439-47. doi:10.1002/hep.20766
 - 27 Varga ZV, Erdelyi K, Paloczi J, et al. Disruption of Renal Arginine Metabolism Promotes Kidney Injury in Hepatorenal Syndrome in Mice. *Hepatology* 2018;68:1519-33. doi:10.1002/hep.29915
 - 28 Thabut D, Massard J, Gangloff A, et al. Model for end-stage liver disease score and systemic inflammatory response are major prognostic factors in patients with cirrhosis and acute functional renal failure. *Hepatology* 2007;46:1872-82. doi:10.1002/hep.21920
 - 29 Solé C, Solà E, Huelin P, et al. Characterization of inflammatory response in hepatorenal syndrome: Relationship with kidney outcome and survival. *Liver Int* 2019;39:1246-55. doi:10.1111/liv.14037
 - 30 Shah N, Mohamed FE, Jover-Cobos M, et al. Increased renal expression and urinary excretion of TLR4 in acute kidney injury associated with cirrhosis. *Liver Int* 2013;33:398-409. doi:10.1111/liv.12047
 - 31 Shah N, Dhar D, El Zahraa Mohammed F, et al. Prevention of acute kidney injury in a rodent model of cirrhosis following selective gut decontamination is associated with reduced renal TLR4 expression. *J Hepatol* 2012;56:1047-53. doi:10.1016/j.jhep.2011.11.024
 - 32 Rivolta R, Maggi A, Cazzaniga M, et al. Reduction of renal cortical blood flow assessed by Doppler in cirrhotic patients with refractory ascites. *Hepatology* 1998;28:1235-40. doi:10.1002/hep.510280510
 - 33 Mindikoglu AL, Dowling TC, Wong-You-Cheong JJ, et al. A pilot study to evaluate renal hemodynamics in cirrhosis by simultaneous glomerular filtration rate, renal plasma flow, renal resistive indices and biomarkers measurements. *Am J Nephrol* 2014;39:543-52. doi:10.1159/000363584
 - 34 Jang HR, Rabb H. Immune cells in experimental acute kidney injury. *Nat Rev Nephrol* 2015;11:88-101. doi:10.1038/nrneph.2014.180
 - 35 Jang JY, Kim TY, Sohn JH, et al. Relative adrenal insufficiency in chronic liver disease: its prevalence and effects on long-term mortality. *Aliment Pharmacol Ther* 2014;40:819-26. doi:10.1111/apt.12891
 - 36 Singh RR, Wallia R, Sachdeva N, Bhalla A, Singh A, Singh V. Relative adrenal insufficiency in cirrhotic patients with ascites (hepatoadrenal syndrome). *Dig Liver Dis* 2018;50:1232-7. doi:10.1016/j.dld.2018.05.011
 - 37 Piano S, Favaretto E, Tonon M, et al. Including Relative Adrenal Insufficiency in Definition and Classification of Acute-on-Chronic Liver Failure. *Clin Gastroenterol Hepatol* 2020;18:1188-1196.e3. doi:10.1016/j.cgh.2019.09.035
 - 38 Acevedo J, Fernández J, Prado V, et al. Relative adrenal insufficiency in decompensated cirrhosis: Relationship to short-term risk of severe sepsis, hepatorenal syndrome, and death. *Hepatology* 2013;58:1757-65. doi:10.1002/hep.26535
 - 39 Tsai MH, Peng YS, Chen YC, et al. Adrenal insufficiency in patients with cirrhosis, severe sepsis and septic shock. *Hepatology* 2006;43:673-81. doi:10.1002/hep.21101
 - 40 Kim G, Huh JH, Lee KJ, Kim MY, Shim KY, Baik SK. Relative Adrenal Insufficiency in Patients with Cirrhosis: A Systematic Review and Meta-Analysis. *Dig Dis Sci* 2017;62:1067-79. doi:10.1007/s10620-017-4471-8
 - 41 Marik PE. Adrenal-exhaustion syndrome in patients with liver disease. *Intensive Care Med* 2006;32:275-80. doi:10.1007/s00134-005-0005-5
 - 42 Bornstein SR. Predisposing factors for adrenal insufficiency. *N Engl J Med* 2009;360:2328-39. doi:10.1056/NEJMra0804635
 - 43 Fernández J, Escorsell A, Zabalza M, et al. Adrenal insufficiency in patients with cirrhosis and septic shock: Effect of treatment with hydrocortisone on survival. *Hepatology* 2006;44:1288-95. doi:10.1002/hep.21352
 - 44 Bräsen JH, Mederacke YS, Schmitz J, et al. Cholemic Nephropathy Causes Acute Kidney Injury and Is Accompanied by Loss of Aquaporin 2 in Collecting Ducts. *Hepatology* 2019;69:2107-19. doi:10.1002/hep.30499
 - 45 van Slambrouck CM, Salem F, Meehan SM, Chang A. Bile cast nephropathy is a common pathologic finding for kidney injury associated with severe liver dysfunction[published Online First: 2013/03/15]. *Kidney Int* 2013;84:192-7. doi:10.1038/ki.2013.78.
 - 46 Nazar A, Pereira GH, Guevara M, et al. Predictors of response to therapy with terlipressin and albumin in patients with cirrhosis and type 1 hepatorenal syndrome. *Hepatology* 2010;51:219-26. doi:10.1002/hep.23283
 - 47 Barreto R, Fagundes C, Guevara M, et al. Type-1 hepatorenal syndrome associated with infections in cirrhosis: natural history, outcome of kidney function, and survival. *Hepatology* 2014;59:1505-13. doi:10.1002/hep.26687
 - 48 Krones E, Eller K, Pollheimer MJ, et al. NorUrsodeoxycholic acid ameliorates cholemic nephropathy in bile duct ligated mice. *J Hepatol* 2017;67:110-9. doi:10.1016/j.jhep.2017.02.019
 - 49 Umgelter A, Reindl W, Wagner KS, et al. Effects of plasma expansion with albumin and paracentesis on haemodynamics and kidney function in critically ill cirrhotic patients with tense ascites and hepatorenal syndrome: a prospective uncontrolled trial. *Crit Care* 2008;12:R4. doi:10.1186/cc6765
 - 50 Velez JCQ, Petkovich B, Karakala N, Huggins JT. Point-of-Care Echocardiography Unveils Misclassification of Acute Kidney Injury as Hepatorenal Syndrome. *Am J Nephrol* 2019;50:204-11. doi:10.1159/000501299
 - 51 Lang F, Tschernko E, Schulze E, et al. Hepatorenal reflex regulating kidney function. *Hepatology* 1991;14:590-4. doi:10.1002/hep.1840140403
 - 52 Ming Z, Smyth DD, Lautt WW. Decreases in portal flow trigger a hepatorenal reflex to inhibit renal sodium and water excretion in rats: role of adenosine. *Hepatology* 2002;35:167-75. doi:10.1053/jhep.2002.30425
 - 53 Garcia-Tsao G, Parikh CR, Viola A. Acute kidney injury in cirrhosis. *Hepatology* 2008;48:2064-77. doi:10.1002/hep.22605
 - 54 Huelin P, Piano S, Solà E, et al. Validation of a Staging System for Acute Kidney Injury in Patients With Cirrhosis and Association With Acute-on-Chronic Liver Failure. *Clin Gastroenterol Hepatol* 2017;15:438-445.e5. doi:10.1016/j.cgh.2016.09.156
 - 55 Dudley FJ, Kanel GC, Wood LJ, Reynolds TB. Hepatorenal syndrome without avid sodium retention. *Hepatology* 1986;6:248-51. doi:10.1002/hep.1840060216
 - 56 Alsaad AA, Wadei HM. Fractional excretion of sodium in hepatorenal syndrome: Clinical and pathological correlation. *World J Hepatol* 2016;8:1497-501. doi:10.4254/wjh.v8.i34.1497
 - 57 Carvounis CP, Nisar S, Guro-Razuman S. Significance of the fractional excretion of urea in the differential diagnosis of acute renal failure. *Kidney Int* 2002;62:2223-9. doi:10.1046/j.1523-1755.2002.00683.x
 - 58 Patidar KR, Kang L, Bajaj JS, Carl D, Sanyal AJ. Fractional excretion of urea: A simple tool for the differential diagnosis of acute kidney injury in cirrhosis. *Hepatology* 2018;68:224-33. doi:10.1002/hep.29772
 - 59 Francoz C, Nadim MK, Durand F. Kidney biomarkers in cirrhosis. *J Hepatol* 2016;65:809-24. doi:10.1016/j.jhep.2016.05.025
 - 60 Allegretti AS, Solà E, Ginès P. Clinical application of kidney biomarkers in cirrhosis. *Am J Kidney Dis* 2020;S0272-6386(20)30691-0.
 - 61 Fagundes C, Pépin MN, Guevara M, et al. Urinary neutrophil gelatinase-associated lipocalin as biomarker in the differential diagnosis of impairment of kidney function in cirrhosis. *J Hepatol* 2012;57:267-73. doi:10.1016/j.jhep.2012.03.015
 - 62 Verna EC, Brown RS, Farrand E, et al. Urinary neutrophil gelatinase-associated lipocalin predicts mortality and identifies acute kidney injury in cirrhosis. *Dig Dis Sci* 2012;57:2362-70. doi:10.1007/s10620-012-2180-x
 - 63 Belcher JM, Sanyal AJ, Peixoto AJ, et al. TRIBE-AKI Consortium. Kidney biomarkers and differential diagnosis of patients with cirrhosis and acute kidney injury. *Hepatology* 2014;60:622-32. doi:10.1002/hep.26980
 - 64 Huelin P, Solà E, Elia C, et al. Neutrophil Gelatinase-Associated Lipocalin for Assessment of Acute Kidney Injury in Cirrhosis: A Prospective Study. *Hepatology* 2019;70:319-33. doi:10.1002/hep.30592
 - 65 Hamdy HS, El-Ray A, Salaheldin M, et al. Urinary Neutrophil Gelatinase-Associated Lipocalin in Cirrhotic Patients with Acute Kidney Injury. *Ann Hepatol* 2018;17:624-30. doi:10.5604/01.3001.0012.0931
 - 66 Barreto R, Elia C, Solà E, et al. Urinary neutrophil gelatinase-associated lipocalin predicts kidney outcome and death in patients with cirrhosis and bacterial infections. *J Hepatol* 2014;61:35-42. doi:10.1016/j.jhep.2014.02.023
 - 67 Ginès A, Escorsell A, Ginès P, et al. Incidence, predictive factors, and prognosis of the hepatorenal syndrome in cirrhosis with ascites. *Gastroenterology* 1993;105:229-36. doi:10.1016/0016-5085(93)90031-7
 - 68 Wong F, Jepsen P, Watson H, Vilstrup H. Un-precipitated acute kidney injury is uncommon among stable patients with cirrhosis and ascites. *Liver Int* 2018;38:1785-92. doi:10.1111/liv.13738
 - 69 Follo A, Llovet JM, Navasa M, et al. Renal impairment after spontaneous bacterial peritonitis in cirrhosis: incidence, clinical course, predictive factors and prognosis. *Hepatology* 1994;20:1495-501. doi:10.1002/hep.1840200619

- 70 Terra C, Guevara M, Torre A, et al. Renal failure in patients with cirrhosis and sepsis unrelated to spontaneous bacterial peritonitis: value of MELD score. *Gastroenterology* 2005;129:1944-53. doi:10.1053/j.gastro.2005.09.024
- 71 Fasolato S, Angeli P, Dallagnese L, et al. Renal failure and bacterial infections in cirrhosis: epidemiology and clinical features. *Hepatology* 2007;45:223-9. doi:10.1002/hep.21443
- 72 Navasa M, Follo A, Filella X, et al. Tumor necrosis factor and interleukin-6 in spontaneous bacterial peritonitis in cirrhosis: relationship with the development of renal impairment and mortality. *Hepatology* 1998;27:1227-32. doi:10.1002/hep.510270507
- 73 Sort P, Navasa M, Arroyo V, et al. Effect of intravenous albumin on renal impairment and mortality in patients with cirrhosis and spontaneous bacterial peritonitis. *N Engl J Med* 1999;341:403-9. doi:10.1056/NEJM199908053410603
- 74 Salerno F, Navickis RJ, Wilkes MM. Albumin infusion improves outcomes of patients with spontaneous bacterial peritonitis: a meta-analysis of randomized trials. *Clin Gastroenterol Hepatol* 2013;11:123-30.e1. doi:10.1016/j.cgh.2012.11.007
- 75 Thévenot T, Bureau C, Oberti F, et al. Effect of albumin in cirrhotic patients with infection other than spontaneous bacterial peritonitis. A randomized trial. *J Hepatol* 2015;62:822-30. doi:10.1016/j.jhep.2014.11.017
- 76 Fernández J, Angeli P, Trebicka J, et al. Efficacy of Albumin Treatment for Patients with Cirrhosis and Infections Unrelated to Spontaneous Bacterial Peritonitis. *Clin Gastroenterol Hepatol* 2020;18:963-973.e14. doi:10.1016/j.cgh.2019.07.055
- 77 Guevara M, Terra C, Nazar A, et al. Albumin for bacterial infections other than spontaneous bacterial peritonitis in cirrhosis. A randomized, controlled study. *J Hepatol* 2012;57:759-65. doi:10.1016/j.jhep.2012.06.013
- 78 Leão GS, John Neto G, Jotz RF, Mattos AA, Mattos AZ. Albumin for cirrhotic patients with extraperitoneal infections: A meta-analysis. *J Gastroenterol Hepatol* 2019;34:2071-6. doi:10.1111/jgh.14791
- 79 Caraceni P, Riggio O, Angeli P, et al, ANSWER Study Investigators. Long-term albumin administration in decompensated cirrhosis (ANSWER): an open-label randomised trial. *Lancet* 2018;391:2417-29. doi:10.1016/S0140-6736(18)30840-7
- 80 Solà E, Solé C, Simón-Talero M, et al. Midodrine and albumin for prevention of complications in patients with cirrhosis awaiting liver transplantation. A randomized placebo-controlled trial. *J Hepatol* 2018;69:1250-9. doi:10.1016/j.jhep.2018.08.006
- 81 Fernández J, Navasa M, Planas R, et al. Primary prophylaxis of spontaneous bacterial peritonitis delays hepatorenal syndrome and improves survival in cirrhosis. *Gastroenterology* 2007;133:818-24. doi:10.1053/j.gastro.2007.06.065
- 82 Kamal F, Khan MA, Khan Z, et al. Rifaximin for the prevention of spontaneous bacterial peritonitis and hepatorenal syndrome in cirrhosis: a systematic review and meta-analysis. *Eur J Gastroenterol Hepatol* 2017;29:1109-17. doi:10.1097/MEG.0000000000000940
- 83 Ruiz-del-Arbol L, Monesillo A, Jimenez W, Garcia-Plaza A, Arroyo V, Rodés J. Paracentesis-induced circulatory dysfunction: mechanism and effect on hepatic hemodynamics in cirrhosis. *Gastroenterology* 1997;113:579-86. doi:10.1053/gast.1997.v113.pm9247479
- 84 Ginès P, Titó L, Arroyo V, et al. Randomized comparative study of therapeutic paracentesis with and without intravenous albumin in cirrhosis. *Gastroenterology* 1988;94:1493-502. doi:10.1016/0016-5085(88)90691-9
- 85 Bernardi M, Caraceni P, Navickis RJ, Wilkes MM. Albumin infusion in patients undergoing large-volume paracentesis: a meta-analysis of randomized trials. *Hepatology* 2012;55:1172-81. doi:10.1002/hep.24786
- 86 Fernández J, Monteagudo J, Bargallo X, et al. A randomized unblinded pilot study comparing albumin versus hydroxyethyl starch in spontaneous bacterial peritonitis. *Hepatology* 2005;42:627-34. doi:10.1002/hep.20829
- 87 Garcia-Martinez R, Caraceni P, Bernardi M, Gines P, Arroyo V, Jalan R. Albumin: pathophysiologic basis of its role in the treatment of cirrhosis and its complications. *Hepatology* 2013;58:1836-46. doi:10.1002/hep.26338
- 88 Zhang WJ, Frei B. Albumin selectively inhibits TNF alpha-induced expression of vascular cell adhesion molecule-1 in human aortic endothelial cells. *Cardiovasc Res* 2002;55:820-9. doi:10.1016/S0008-6363(02)00492-3
- 89 Chen TA, Tsao YC, Chen A, et al. Effect of intravenous albumin on endotoxin removal, cytokines, and nitric oxide production in patients with cirrhosis and spontaneous bacterial peritonitis. *Scand J Gastroenterol* 2009;44:619-25. doi:10.1080/0036520902719273
- 90 China L, Maini A, Skene SS, et al. Albumin Counteracts Immune-Suppressive Effects of Lipid Mediators in Patients With Advanced Liver Disease. *Clin Gastroenterol Hepatol* 2018;16:738-747.e7. doi:10.1016/j.cgh.2017.08.027
- 91 China L, Skene SS, Bennett K, et al. ATTIRE: Albumin To prevent Infection in chronic liver failure: study protocol for an interventional randomised controlled trial. *BMJ Open* 2018;8:e023754.
- 92 European Association for the Study of the Liver. Electronic address: easloffice@easloffice.eu, European Association for the Study of the Liver. EASL Clinical Practice Guidelines for the management of patients with decompensated cirrhosis. *J Hepatol* 2018;69:406-60. doi:10.1016/j.jhep.2018.03.024
- 93 Bhutta AQ, Garcia-Tsao G, Reddy KR, et al. Beta-blockers in hospitalised patients with cirrhosis and ascites: mortality and factors determining discontinuation and reinitiation. *Aliment Pharmacol Ther* 2018;47:78-85. doi:10.1111/apt.14366
- 94 Mandorfer M, Bota S, Schwabl P, et al. Nonselective β blockers increase risk for hepatorenal syndrome and death in patients with cirrhosis and spontaneous bacterial peritonitis. *Gastroenterology* 2014;146:1680-90.e1. doi:10.1053/j.gastro.2014.03.005
- 95 Ibrahim ES, Alsebaey A, Zagha H, Moawad Abdelmageed S, Gameel K, Abdelsameea E. Long-term rifaximin therapy as a primary prevention of hepatorenal syndrome. *Eur J Gastroenterol Hepatol* 2017;29:1247-50. doi:10.1097/MEG.0000000000000967
- 96 Dong T, Aronsohn A, Gautham Reddy K, Te HS. Rifaximin Decreases the Incidence and Severity of Acute Kidney Injury and Hepatorenal Syndrome in Cirrhosis. *Dig Dis Sci* 2016;61:3621-6. doi:10.1007/s10620-016-4313-0
- 97 Velez JC, Nietert PJ. Therapeutic response to vasoconstrictors in hepatorenal syndrome parallels increase in mean arterial pressure: a pooled analysis of clinical trials. *Am J Kidney Dis* 2011;58:928-38. doi:10.1053/j.ajkd.2011.07.017
- 98 Facciorusso A, Chandar AK, Murad MH, et al. Comparative efficacy of pharmacological strategies for management of type 1 hepatorenal syndrome: a systematic review and network meta-analysis. *Lancet Gastroenterol Hepatol* 2017;2:94-102. doi:10.1016/S2468-1253(16)30157-1
- 99 Jamil K, Pappas SC, Devarakonda KR. In vitro binding and receptor-mediated activity of terlipressin at vasopressin receptors V_1 and V_2 . *J Exp Pharmacol* 2017;10:1-7. doi:10.2147/JEP.S146034
- 100 Tanoue A, Ito S, Honda K, et al. The vasopressin V_{1b} receptor critically regulates hypothalamic-pituitary-adrenal axis activity under both stress and resting conditions. *J Clin Invest* 2004;113:302-9. doi:10.1172/JCI200419656
- 101 Wong F, Pappas SC, Boyer TD, et al, REVERSE Investigators. Terlipressin Improves Renal Function and Reverses Hepatorenal Syndrome in Patients With Systemic Inflammatory Response Syndrome. *Clin Gastroenterol Hepatol* 2017;15:266-272.e1. doi:10.1016/j.cgh.2016.07.016
- 102 Russell JA, Walley KR. Vasopressin and its immune effects in septic shock. *J Innate Immun* 2010;2:446-60. doi:10.1159/000318531
- 103 Krag A, Bendtsen F, Pedersen EB, Holstein-Rathlou NH, Møller S. Effects of terlipressin on the aquaretic system: evidence of antidiuretic effects. *Am J Physiol Renal Physiol* 2008;295:F1295-300. doi:10.1152/ajprenal.90407.2008
- 104 Solà E, Lens S, Guevara M, et al. Hyponatremia in patients treated with terlipressin for severe gastrointestinal bleeding due to portal hypertension. *Hepatology* 2010;52:1783-90. doi:10.1002/hep.23893
- 105 Laterre PF, Berry SM, Blemings A, et al, SEPSIS-ACT Investigators. Effect of Selepressin vs Placebo on Ventilator- and Vasopressor-Free Days in Patients With Septic Shock: The SEPSIS-ACT Randomized Clinical Trial. *JAMA* 2019;322:1476-85. doi:10.1001/jama.2019.14607
- 106 He X, Su F, Taccone FS, et al. A Selective V_{1A} Receptor Agonist, Selepressin, Is Superior to Arginine Vasopressin and to Norepinephrine in Ovine Septic Shock. *Crit Care Med* 2016;44:23-31. doi:10.1097/CCM.0000000000001380
- 107 Merkel C, Gatta A, Zuin R, Finucci GF, Nosadini R, Ruol A. Effect of somatostatin on splanchnic hemodynamics in patients with liver cirrhosis and portal hypertension. *Digestion* 1985;32:92-8. doi:10.1159/000199224
- 108 Møller S, Brinch K, Henriksen JH, Becker U. Effect of octreotide on systemic, central, and splanchnic haemodynamics in cirrhosis. *J Hepatol* 1997;26:1026-33. doi:10.1016/S0168-8278(97)80111-0
- 109 Pomier-Layrargues G, Paquin SC, Hassoun Z, Laforce M, Tran A. Octreotide in hepatorenal syndrome: a randomized, double-blind, placebo-controlled, crossover study. *Hepatology* 2003;38:238-43. doi:10.1053/jhep.2003.50276
- 110 Angeli P, Volpin R, Piovon D, et al. Acute effects of the oral administration of midodrine, an alpha-adrenergic agonist, on renal hemodynamics and renal function in cirrhotic patients with ascites. *Hepatology* 1998;28:937-43. doi:10.1002/hep.510280407
- 111 Angeli P, Volpin R, Gerunda G, et al. Reversal of type 1 hepatorenal syndrome with the administration of midodrine and octreotide. *Hepatology* 1999;29:1690-7. doi:10.1002/hep.510290629

- 112 Wong F, Pantea L, Sniderman K. Midodrine, octreotide, albumin, and TIPS in selected patients with cirrhosis and type 1 hepatorenal syndrome. *Hepatology* 2004;40:55-64. doi:10.1002/hep.20262
- 113 Solanki P, Chawla A, Garg R, Gupta R, Jain M, Sarin SK. Beneficial effects of terlipressin in hepatorenal syndrome: a prospective, randomized placebo-controlled clinical trial. *J Gastroenterol Hepatol* 2003;18:152-6. doi:10.1046/j.1440-1746.2003.02934.x
- 114 Neri S, Pulvirenti D, Malaguarnera M, et al. Terlipressin and albumin in patients with cirrhosis and type I hepatorenal syndrome. *Dig Dis Sci* 2008;53:830-5. doi:10.1007/s10620-007-9919-9
- 115 Sanyal AJ, Boyer T, Garcia-Tsao G, et al. Terlipressin Study Group. A randomized, prospective, double-blind, placebo-controlled trial of terlipressin for type 1 hepatorenal syndrome. *Gastroenterology* 2008;134:1360-8. doi:10.1053/j.gastro.2008.02.014
- 116 Boyer TD, Sanyal AJ, Wong F, et al. REVERSE Study Investigators. Terlipressin Plus Albumin Is More Effective Than Albumin Alone in Improving Renal Function in Patients With Cirrhosis and Hepatorenal Syndrome Type 1. *Gastroenterology* 2016;150:1579-1589.e2. doi:10.1053/j.gastro.2016.02.026
- 117 Wong F, Curry M, Reddy R, et al. CONFIRM Study Investigators. LO5 The CONFIRM study: A North American randomized controlled trial (RCT) of terlipressin plus albumin for the treatment of hepatorenal syndrome type 1 (HRS-1). *Hepatology* 2019;70(Suppl 1):10.
- 118 Alessandria C, Ottobrelli A, Debernardi-Venon W, et al. Noradrenalin vs terlipressin in patients with hepatorenal syndrome: a prospective, randomized, unblinded, pilot study. *J Hepatol* 2007;47:499-505. doi:10.1016/j.jhep.2007.04.010
- 119 Sharma P, Kumar A, Shrama BC, Sarin SK. An open label, pilot, randomized controlled trial of noradrenaline versus terlipressin in the treatment of type 1 hepatorenal syndrome and predictors of response. *Am J Gastroenterol* 2008;103:1689-97. doi:10.1111/j.1572-0241.2008.01828.x
- 120 Singh V, Ghosh S, Singh B, et al. Noradrenaline vs. terlipressin in the treatment of hepatorenal syndrome: a randomized study. *J Hepatol* 2012;56:1293-8.
- 121 Cavallin M, Kamath PS, Merli M, et al. Italian Association for the Study of the Liver Study Group on Hepatorenal Syndrome. Terlipressin plus albumin versus midodrine and octreotide plus albumin in the treatment of hepatorenal syndrome: A randomized trial. *Hepatology* 2015;62:567-74. doi:10.1002/hep.27709
- 122 Arora V, Maiwall R, Rajan V, et al. Terlipressin Is Superior to Noradrenaline in the Management of Acute Kidney Injury in Acute on Chronic Liver Failure. *Hepatology* 2020;71:600-10. doi:10.1002/hep.30208
- 123 Ortega R, Ginès P, Uriz J, et al. Terlipressin therapy with and without albumin for patients with hepatorenal syndrome: results of a prospective, nonrandomized study. *Hepatology* 2002;36:941-8. doi:10.1053/jhep.2002.35819
- 124 Cavallin M, Piano S, Romano A, et al. Terlipressin given by continuous intravenous infusion versus intravenous boluses in the treatment of hepatorenal syndrome: A randomized controlled study. *Hepatology* 2016;63:983-92. doi:10.1002/hep.28396
- 125 Duvoux C, Zanditenas D, Hézode C, et al. Effects of noradrenalin and albumin in patients with type I hepatorenal syndrome: a pilot study. *Hepatology* 2002;36:374-80. doi:10.1053/jhep.2002.34343
- 126 Rodríguez E, Elia C, Solà E, et al. Terlipressin and albumin for type-1 hepatorenal syndrome associated with sepsis. *J Hepatol* 2014;60:955-61. doi:10.1016/j.jhep.2013.12.032
- 127 Piano S, Schmidt HH, Ariza X, et al. EASL CLIF Consortium, European Foundation for the Study of Chronic Liver Failure (EF CLIF). Association Between Grade of Acute on Chronic Liver Failure and Response to Terlipressin and Albumin in Patients With Hepatorenal Syndrome. *Clin Gastroenterol Hepatol* 2018;16:1792-1800.e3. doi:10.1016/j.cgh.2018.01.035
- 128 Bortoluzzi A, Ceolotto G, Gola E, et al. Positive cardiac inotropic effect of albumin infusion in rodents with cirrhosis and ascites: molecular mechanisms. *Hepatology* 2013;57:266-76. doi:10.1002/hep.26021
- 129 Stocker R, Glazer AN, Ames BN. Antioxidant activity of albumin-bound bilirubin. *Proc Natl Acad Sci U S A* 1987;84:5918-22. doi:10.1073/pnas.84.16.5918
- 130 Cantin AM, Paquette B, Richter M, Larivée P. Albumin-mediated regulation of cellular glutathione and nuclear factor kappa B activation. *Am J Respir Crit Care Med* 2000;162:1539-46. doi:10.1164/ajrccm.162.4.9910106
- 131 Lejon S, Frick IM, Björck L, Wikström M, Svensson S. Crystal structure and biological implications of a bacterial albumin binding module in complex with human serum albumin. *J Biol Chem* 2004;279:42924-8. doi:10.1074/jbc.M406957200
- 132 Mitzner SR, Stange J, Klammt S, et al. Improvement of hepatorenal syndrome with extracorporeal albumin dialysis MARS: results of a prospective, randomized, controlled clinical trial. *Liver Transpl* 2000;6:277-86. doi:10.1053/lt.2000.6355
- 133 Wong F, Raina N, Richardson R. Molecular adsorbent recirculating system is ineffective in the management of type 1 hepatorenal syndrome in patients with cirrhosis with ascites who have failed vasoconstrictor treatment. *Gut* 2010;59:381-6.
- 134 Ginès P, Uriz J, Calahorra B, et al. Transjugular intrahepatic portosystemic shunting versus paracentesis plus albumin for refractory ascites in cirrhosis. *Gastroenterology* 2002;123:1839-47. doi:10.1053/gast.2002.37073
- 135 Azoulay D, Castaing D, Majno P, et al. Salvage transjugular intrahepatic portosystemic shunt for uncontrolled variceal bleeding in patients with decompensated cirrhosis. *J Hepatol* 2001;35:590-7. doi:10.1016/S0168-8278(01)00185-4
- 136 Guevara M, Ginès P, Bandi JC, et al. Transjugular intrahepatic portosystemic shunt in hepatorenal syndrome: effects on renal function and vasoactive systems. *Hepatology* 1998;28:416-22. doi:10.1002/hep.510280219
- 137 Song T, Rössle M, He F, Liu F, Guo X, Qi X. Transjugular intrahepatic portosystemic shunt for hepatorenal syndrome: A systematic review and meta-analysis. *Dig Liver Dis* 2018;50:323-30. doi:10.1016/j.dld.2018.01.123
- 138 Zhang Z, Maddukuri G, Jaipaul N, Cai CX. Role of renal replacement therapy in patients with type 1 hepatorenal syndrome receiving combination treatment of vasoconstrictor plus albumin. *J Crit Care* 2015;30:969-74. doi:10.1016/j.jccr.2015.05.006
- 139 Lenhart A, Hussain S, Salgia R. Chances of Renal Recovery or Liver Transplantation After Hospitalization for Alcoholic Liver Disease Requiring Dialysis. *Dig Dis Sci* 2018;63:2800-9. doi:10.1007/s10620-018-5170-9
- 140 Fraley DS, Burr R, Bernardini J, Angus D, Kramer DJ, Johnson JP. Impact of acute renal failure on mortality in end-stage liver disease with or without transplantation. *Kidney Int* 1998;54:518-24. doi:10.1046/j.1523-1755.1998.00004.x
- 141 Wong LP, Blackley MP, Andreoni KA, Chin H, Falk RJ, Klemmer PJ. Survival of liver transplant candidates with acute renal failure receiving renal replacement therapy. *Kidney Int* 2005;68:362-70. doi:10.1111/j.1523-1755.2005.00408.x
- 142 Allegretti AS, Parada XV, Eneanya ND, et al. Prognosis of Patients with Cirrhosis and AKI Who Initiate RRT. *Clin J Am Soc Nephrol* 2018;13:16-25. doi:10.2215/CJN.03610417
- 143 European Association for the Study of the Liver. EASL clinical practice guidelines on the management of ascites, spontaneous bacterial peritonitis, and hepatorenal syndrome in cirrhosis. *J Hepatol* 2010;53:397-417. doi:10.1016/j.jhep.2010.05.004
- 144 Iwatsuki S, Popovtzer MM, Corman JL, et al. Recovery from "hepatorenal syndrome" after orthotopic liver transplantation. *N Engl J Med* 1973;289:1155-9. doi:10.1056/NEJM197311292892201
- 145 Runyon BAAASLD Practice Guidelines Committee. Management of adult patients with ascites due to cirrhosis: an update. *Hepatology* 2009;49:2087-107. doi:10.1002/hep.22853
- 146 Israni AK, Xiong H, Liu J, et al. Predicting end-stage renal disease after liver transplant. *Am J Transplant* 2013;13:1782-92. doi:10.1111/ajt.12257
- 147 Organ Procurement and Transplantation Network. Policies. 2020. https://optn.transplant.hrsa.gov/media/1200/optn_policies.pdf.
- 148 Chauhan K, Azzi Y, Faddoul G, et al. Pre-liver transplant renal dysfunction and association with post-transplant end-stage renal disease: A single-center examination of updated UNOS recommendations. *Clin Transplant* 2018;32:e13428. doi:10.1111/ctr.13428
- 149 Nadim MK, Genyk YS, Tokin C, et al. Impact of the etiology of acute kidney injury on outcomes following liver transplantation: acute tubular necrosis versus hepatorenal syndrome. *Liver Transpl* 2012;18:539-48. doi:10.1002/lt.23384
- 150 OPTN/UNOS Kidney Transplantation Committee. Simultaneous Liver Kidney (SLK) Allocation Policy. https://optn.transplant.hrsa.gov/media/1192/0815-12_SLK_Allocation.pdf.
- 151 Runyon BA. Management of adult patients with ascites due to cirrhosis: update 2012. https://www.aasld.org/sites/default/files/2019-06/141020_Guideline_Ascites_4UFb_2015.pdf.
- 152 Conrad KP. Unveiling the vasodilatory actions and mechanisms of relaxin. *Hypertension* 2010;56:2-9. doi:10.1161/HYPERTENSIONAHA.109.133926
- 153 Fallowfield JA, Hayden AL, Snowdon VK, et al. Relaxin modulates human and rat hepatic myofibroblast function and ameliorates portal hypertension in vivo. *Hepatology* 2014;59:1492-504. doi:10.1002/hep.26627
- 154 Snowdon VK, Lachlan NJ, Hoy AM, et al. Serelaxin as a potential treatment for renal dysfunction in cirrhosis: Preclinical evaluation and results of a randomized phase 2 trial. *PLoS Med* 2017;14:e1002248. doi:10.1371/journal.pmed.1002248
- 155 Stine JG, Wang J, Cornella SL, et al. Treatment of Type-1 Hepatorenal Syndrome with Pentoxifylline: A Randomized Placebo Controlled Clinical Trial. *Ann Hepatol* 2018;17:300-6. doi:10.5604/01.3001.0010.8661